

Bosonic stimulation of cold 1s excitons into a harmonic potential minimum in Cu_2O

N. Naka and N. Nagasawa

*Department of Physics, Graduate School of Science, University of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan*

(Dated: 17 December, 2002)

Density distribution of cold exciton clouds generated into a strain-induced potential well by two-photon excitation in Cu_2O is studied at 2 K. We find that an anomalous spike, which can be interpreted as accumulation of the excitons into the ground state, emerges at the potential minimum. The accumulation can be due to stimulated scattering of cold excitons, mediated by acoustic phonon emission. Possibility of the formation of the thermodynamic Bose-Einstein condensate of paraexcitons has been discussed. PACS numbers: 71.35.Lk, 78.35.-y

Before the first realization of atomic Bose-Einstein condensation (BEC) in 1995 [1], excitons in semiconductors were thought to be one of the most favorable candidates for realizing BEC [2]. It has long been believed that excitons in Cu_2O is the best system to pursue such a phase transition in a three-dimensional system, because of their long lifetime derived from the dipole forbidden gap; the orthoexcitons couple with photons only via quadrupole interaction, and paraexcitons are basically inactive for optical transitions, thereby leading to a long lifetime.

However, the long lifetime of the excitons in Cu_2O allows them to expand in free crystals, and means that their recombination rate is low. The former reduces the exciton density against the BEC, and the latter makes it difficult to monitor exciton densities by optical means. In order to overcome these unfavorable situations, we employed inhomogeneous strain that creates potential wells and accelerates the direct recombination rate. We have been studying kinetics of a cold exciton system in a potential well, by using two-photon excitation. In this regime, orthoexcitons are created near the local momentum zero [3, 4]. In this communication, we report on a novel phenomenon showing stimulated scattering of excitons; this can be mediated by acoustic phonon emission.

A natural Cu_2O crystal cut into a 3 mm-cube was pushed against a glass lens of curvature $R = 7.78$ mm to create harmonic potential wells (see the inset of Fig. 1). Solid lines in Fig. 1 show calculated potential shapes for the four lowest exciton states in Cu_2O , i.e. upper-, middle- and lower-branch orthoexcitons and paraexcitons. It is evident that middle- and lower-branch orthoexcitons as well as the paraexcitons become trapped states while the upper-branch orthoexcitons form an untrapped state. The trapping potential shapes are approximately harmonic around the center of the wells. An infrared laser beam from a color-center laser (Solar, LF151) pumped by a Q-switched YAG laser (SantaFe Laser, C-140) was used to inject orthoexcitons into a potential well. The repetition rate and the duration of the laser light were 400 Hz and 12 ns, respectively. The two-photon excitation energy of the laser light was tuned slightly below the orthoexciton resonance under zero stress, so that middle- and lower-branch orthoexcitons were generated into the trap [8]. The luminescence was viewed along the stress axis through a prism mounted




FIG. 1: Solid lines: calculated potential energies of the four 1s exciton branches, i.e. upper-, middle-, and lower-branch orthoexcitons, and paraexcitons. The zero of the energy was taken to be the orthoexciton resonance under zero stress, 2032.78 meV. The inset shows schematic for the experimental configuration. Two potential minima were created near the center of the contact to a glass plunger, while data for positive y are shown. The shadowed profile shows spatial distribution of the exciton cloud, measured at $t = 10$ ns. Dashed arrows labeled 1 and 2 show the proposed mechanism of stimulated scattering after downconversion of orthoexcitons to paraexcitons at the rim (see the text for detail).

below the crystal. A $\times 20$ magnified image of the luminous cloud was detected by an ICCD camera system (LaVision, Picostar HR12) after being passed through a filter which cuts off the infrared scattering light. By inserting a spectrometer before the ICCD camera system, we confirmed that luminescence light consists of intrin-




FIG. 2: Time evolution of the spatial profile of the exciton cloud pumped at a rim of a potential well. The baseline for each profile has been shifted for clarity. A sharp spike emerges at $y \simeq 300 \mu\text{m}$, and rapidly grows after $t = 4 \text{ ns}$. Part (b) shows profiles after $t = 10 \text{ ns}$. The gate width was set at 4 ns, and the power of the incident light was $I = 9 \text{ mW}$ on the sample surface.

sic free-exciton recombination luminescence. The spatial resolution of the optical system was higher than $10 \mu\text{m}$.

Figure 2 shows time-resolved profile of the exciton clouds at different times, t , after the incidence of the laser light. The gate width was set at 4 ns, and respective profiles were extracted from luminescence images averaged over 4000 laser shots. Parts (a) and (b) show growing and decaying of the exciton density, with time interval of 2 ns. Three major peaks are seen in a range between $y = 200 \mu\text{m}$ and $y = 400 \mu\text{m}$. The intensity of the side peaks was almost proportional to the temporal profile of the laser pulse, and to the square of the power density of the laser light. On the other hand, the central peak shows rapid growth even after the maximum of the incident laser intensity at $t = 6 \text{ ns}$, and has nonlinear dependence on the incident power density. The width of the central peak was $7.7 \mu\text{m}$, much narrower than that of side peaks.

In order to understand the origin of the side peaks and of the anomalous spike in between, the spatial profile at $t = 10 \text{ ns}$ is superimposed to the potential shapes in Fig. 1. The position of the baseline represents the two-photon excitation energy, measured from the orthoexciton resonance under zero stress, 2032.78 meV . Resonance creation of orthoexcitons occurs in selected regions where the two-photon excitation energy coincides with the local exciton energy; structures marked L, M and L', M' are due to the resonance with lower- and middle-branch

orthoexcitons at a rim of the trap. On the other hand, emergence of the sharp spike, C, at the center of the well cannot be explained by resonance to any orthoexciton branches. This difference, from the case for side peaks, is consistent with the above features, i.e. nonlinearity and the delay in the growing process for the central spike, C.

The position of the center of the well for middle-branch orthoexcitons is almost the same as that for paraexcitons. Therefore, judging from its position, the origin of the spike can be middle-branch orthoexcitons, paraexcitons, or both. Unfortunately, however, we could not obtain spectrally resolved data in the present experiment, because of the weakness of the signal. Dashed arrows in Fig. 1 show a possible mechanism for the emergence of the spike: the orthoexciton-paraexciton downconversion occurs in the injection spot at a rim (the arrow labeled 1), and then paraexcitons at the rim are scattered into the center of the well (the arrow labeled 2). The energy transfer necessary for this scattering process is inferred to be 0.6 meV , in agreement with the energy position where the acoustic (LA)-phonon dispersion crosses the exciton dispersion. Therefore, the scattering can be mediated by acoustic phonon emission. In fact, we found that the central spike appears only when the laser beam traverses the center of the well, and when the two-photon excitation energy is set at $\Delta E = -5.03 \pm 0.25 \text{ meV}$. Considering the timescale of the downconversion rate, 3 ns [5], and the energy transfer, 1.0 meV or more, necessary for the scattering of orthoexcitons from the rim to their potential minimum, condensation of the middle-branch orthoexcitons is not likely. To conclude, the spike marked C is most likely to be due to paraexcitons accumulated in the ground state at the center of the potential well.

The role of the acoustic phonon in the accumulation of the condensate has been discussed in Refs. [6, 7]. The building time has been estimated as a few nanoseconds. The observed delay is in quantitative agreement with this prediction, provided that the central spike is due to a condensate. Using the measured size of the central spike, the phase-space density of paraexcitons was estimated as ≥ 1 . A more detailed measurement of the building time of the spike deserves future study. Confirmation with higher spectral resolution, which enables measurements of the size of the ground-state wavefunction, is in progress.

We would like to thank Mr. T. Ueda (Mrubun Co.) and Prof. M. Kuwata-Gonokami (Univ. Tokyo) for their kind arrangement for using the ICCD camera system, and Mr. T. Nishi and Mr. T. Mizuno (Marubun Co.) for their technical support. This work was partially supported by The Mitsubishi foundation and the grant-in-aid for scientific research from the Ministry of Education, Science and Culture, Japan.

[1] M.H. Anderson, J.R. Ensher, M.R. Matthews, C.E. Wiemann, and E.A. Cornell, *Science* **269** (1995) 198.

[2] *Bose-Einstein Condensation*, A. Griffin, D.W. Snoke, and S. Stringari, eds., (Cambridge University Press, Cam-

- bridge, 1995).
- [3] N. Naka and N. Nagasawa, *Phys. Rev. B* **65** (2002) 075209.
 - [4] N. Naka and N. Nagasawa, *Phys. Rev. B* **65** (2002) 245203.
 - [5] J.S. Weiner, N. Caswell, P.Y. Yu, and A. Mysyrowicz, *Solid State Commun.* **46** (1983) 105.
 - [6] O.M. Schmitt, D.B. Tran Thoai, L. Bányai, P. Gartner, and H. Haug, *Phys. Rev. Lett.* **86** (2001) 3839.
 - [7] A. Schmitt, L. Bányai, and H. Haug, *Phys. Rev. B* **63** (2001) 205113.
 - [8] The two-photon excitation of paraexcitons is basically possible, but is not practical to generate a high-density gas of excitons [4].